arXiv:1209.4604v2 [math.NT] 24 Oct 2012

SOME PROPERTIES OF FIBONACCI NUMBERS

ALEXANDRE LAUGIER AND MANJIL P. SAIKIA

Abstract. In this paper we study the Fibonacci numbers and derive some interesting properties and recurrence relations. We also define period of a Fibonacci sequence modulo an integer, m and derive certain interesting properties related to them.

1. Preliminaries

We begin with the following famous results without proof.

Lemma 1.1 (Euclid)**.** *If* $ab \equiv 0 \pmod{p}$ *with* a, b *two integers and* p *a prime, then either* $p|a$ *or* $p|b$.

Remark 1.2. *In particular, if* $gcd(a, b) = 1$, p *divides only one of the numbers* a, b.

Theorem 1.3 (Fermat's Little Theorem)**.** *If* p *is a prime and* $n \in \mathbb{N}$ *relatively prime to* p, *then* $n^{p-1} \equiv 1 \pmod{p}$.

Theorem 1.4 (Euler)**.** *Let* p *be an odd prime and* $\gcd(a, p) = 1$. *Then* a *is a quadratic nonresidue of* p *if* $a^{\frac{p-1}{2}} \equiv -1 \pmod{p}$.

Definition 1.5. *Let* p *be an odd prime and let* $\gcd(a, p) = 1$. *The Legendre symbol* (a/p) *is defined to be equal to* 1 *if* a *is a quadratic residue of* p *and is equal to* -1 *is* a *is a quadratic non residue of* p.

Theorem 1.6.

$$(5/5k + 2) = -1.$$

Proof. Clearly $(5/5k + 2) = (5k + 2/5)$ since $5 \equiv 1 \pmod{4}$.

Again $(5k + 2/5) = (2/5)$ since $5k + 2 \equiv 2 \pmod{5}$.

Also it is a well known fact that $(2/5) = -1$ since $5 \equiv 5 \pmod{8}$. □

For proofs of the above theorems the reader is suggested to see [1] or [2].

Theorem 1.7. *If* $x^2 \equiv 1 \pmod{p}$ *with* p *a prime, then either* $x \equiv 1 \pmod{p}$ *or* $x \equiv p - 1 \pmod{p}$.

Proof. If $x^2 \equiv 1 \pmod{p}$ with p a prime, then we have

$$x^2 - 1 \equiv 0 \pmod{p}$$

$$(x - 1)(x + 1) \equiv 0 \pmod{p}$$

$x - 1 \equiv 0 \pmod{p}$ or $x + 1 \equiv 0 \pmod{p}$. It is equivalent to say that $x \equiv 1 \pmod{p}$ or $x \equiv -1 \equiv p - 1 \pmod{p}$. □

Second author is supported by DST-INSPIRE Scholarship 422/2009.

Let p a prime number such that $p = 5k + 2$ with k an odd positive integer. From Fermat's Little Theorem we have

$$\left(5^{\frac{5k+1}{2}}\right)^2 \equiv 1 \pmod{5k+2}.$$

From Theorem 1.7, we have either $5^{\frac{5k+1}{2}} \equiv 1 \pmod{5k+2}$ or $5^{\frac{5k+1}{2}} \equiv 5k+1 \pmod{5k+2}$. Moreover, we can observe that

$$5(2k+1) \equiv 1 \pmod{5k+2}.$$

Theorem 1.8.

$$5^{\frac{5k+1}{2}} \equiv 5k+1 \pmod{5k+2}$$

where $5k + 2$ *is a prime.*

The proof of Theorem 1.8 follows very easily from Theorems 1.4 and 1.6.

We fix the notation $[[1, n]] = \{1, 2, \dots, n\}$ throughout the rest of the paper. We now have the following properties.

Property 1.9.

$$\binom{5k+1}{2l+1} \equiv 5k+1 \pmod{5k+2},$$

with $l \in [[0, \lfloor \frac{5k}{2} \rfloor]]$ *and* $5k + 2$ *is a prime.*

Proof. Notice that for $l = 0$ the property is obviously true.

We also have

$$\binom{5k+1}{2l+1} = \frac{(5k+1)5k(5k-1)\dots(5k-2l+1)}{(2l+1)!}.$$

Or,

$$\begin{aligned} 5k &\equiv -2 \pmod{5k+2}, \\ 5k-1 &\equiv -3 \pmod{5k+2}, \\ &\vdots \\ 5k-2l+1 &\equiv -(2l+1) \pmod{5k+2}. \end{aligned}$$

Multiplying these congruences we get

$$5k(5k-1)\dots(5k-2l+1) \equiv (2l+1)! \pmod{5k+2}.$$

Therfore

$$(2l+1)!\binom{5k+1}{2l+1} \equiv (5k+1)(2l+1)! \pmod{5k+2}.$$

Since $(2l+1)!$ and $5k+2$ are relatively prime, we obtain

$$\binom{5k+1}{2l+1} \equiv 5k+1 \pmod{5k+2}.$$

□

Property 1.10.

$$\binom{5k}{2l+1} \equiv 5k-2l \equiv -2(l+1) \pmod{5k+2}$$

with $l \in [[0, \lfloor \frac{5k}{2} \rfloor]]$ *and* $5k + 2$ *is a prime.*

Proof. Notice that for $l = 0$ the property is obviously true.

We have

$$\binom{5k}{2l+1} = \frac{5k(5k-1)\dots(5k-2l+1)(5k-2l)}{(2l+1)!}.$$

Or

$$\begin{gathered} 5k \equiv -2 \pmod{5k+2}, \\ 5k-1 \equiv -3 \pmod{5k+2}, \\ \vdots \\ 5k-2l+1 \equiv -(2l+1) \pmod{5k+2}. \end{gathered}$$

Multiplying these congruences we get

$$5k(5k-1)\dots(5k-2l+1) \equiv (2l+1)! \pmod{5k+2}.$$

Therfore

$$(2l+1)!\binom{5k}{2l+1} \equiv (2l+1)!(5k-2l) \pmod{5k+2}.$$

Since $(2l+1)!$ and $5k+2$ are relatively prime, we obtain

$$\binom{5k+1}{2l+1} \equiv 5k-2l \equiv 5k+2-2-2l \equiv -2(l+1) \pmod{5k+2}.$$

$\square$

2. Formulas for the Fibonacci numbers

The Fibonacci sequence (F_n) is defined by $F_0 = 0, F_1 = 1, F_{n+2} = F_n + F_{n+1}$.

The first Fibonacci numbers are

$F_0 = 0$, $F_1 = 1$, $F_2 = 1$, $F_3 = 2$, $F_4 = 3$, $F_5 = 5$, $F_6 = 8$, $F_7 = 13$, $F_8 = 21$, $F_9 = 34$, $F_{10} = 55$,

$F_{11} = 89$, $F_{12} = 144$, $F_{13} = 233$, $F_{14} = 377$, $F_{15} = 610$, $F_{16} = 987$, $F_{17} = 1597$, $F_{18} = 2584, \dots$

From the definition of the Fibonacci sequences it can be established the formula for the *nth* Fibonacci number

$$F_n = \frac{\varphi^n - (1-\varphi)^n}{\sqrt{5}}$$

where $\varphi = \frac{1+\sqrt{5}}{2}$ is the golden ratio.

From binomial theorem, we have for $a \neq 0$ and $n \in \mathbb{N}$

$$(a+b)^n - (a-b)^n = \sum_{k=0}^{n}\binom{n}{k}a^{n-k}b^k(1-(-1)^k) = 2\sum_{l=0}^{\lfloor\frac{n-1}{2}\rfloor}\binom{n}{2l+1}a^{n-(2l+1)}b^{2l+1},$$

$$(a+b)^n - (a-b)^n = 2a^n\sum_{l=0}^{\lfloor\frac{n-1}{2}\rfloor}\binom{n}{2l+1}\left(\frac{b}{a}\right)^{2l+1}. \tag{2.1}$$

We set

$$\begin{gathered} a+b = \varphi = \tfrac{1+\sqrt{5}}{2}, \\ a-b = 1-\varphi = \tfrac{1-\sqrt{5}}{2}. \end{gathered}$$

So

$$a = \frac{1}{2},\ b = \frac{2\varphi-1}{2} = \frac{\sqrt{5}}{2}.$$

Thus

$$\frac{b}{a} = \sqrt{5}.$$

We get from (2.1)

$$\varphi^n - (1-\varphi)^n = \frac{\sqrt{5}}{2^{n-1}} \sum_{l=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n}{2l+1} 5^l.$$

Thus we have

Theorem 2.1.

$$F_n = \frac{1}{2^{n-1}} \sum_{l=0}^{\lfloor \frac{n-1}{2} \rfloor} \binom{n}{2l+1} 5^l.$$

For instance,

$$n = 1: \; F_1 = \frac{1}{2^0} \sum_{l=0}^{0} \binom{1}{2l+1} 5^l = 1,$$

$$n = 2: \; F_2 = \frac{1}{2} \sum_{l=0}^{0} \binom{2}{2l+1} 5^l = 1,$$

$$n = 3: \; F_3 = \frac{1}{2^2} \sum_{l=0}^{1} \binom{3}{2l+1} 5^l = \frac{1}{4} \left\{ \binom{3}{1} + \binom{3}{3} \times 5 \right\} = \frac{1}{4}(3+5) = \frac{8}{4} = 2,$$

$$n = 4: \; F_4 = \frac{1}{2^3} \sum_{l=0}^{1} \binom{4}{2l+1} 5^l = \frac{1}{8} \left\{ \binom{4}{1} + \binom{4}{3} \times 5 \right\} = \frac{1}{8}(4+20) = \frac{24}{8} = 3,$$

$$n = 5: \; F_5 = \frac{1}{2^4} \sum_{l=0}^{2} \binom{5}{2l+1} 5^l = \frac{1}{16} \left\{ \binom{5}{1} + \binom{5}{3} \times 5 + \binom{5}{5} \times 5^2 \right\} = \frac{1}{16}(5+50+25) = \frac{80}{16} = 5,$$

$$n = 6: \; F_6 = \frac{1}{2^5} \sum_{l=0}^{2} \binom{6}{2l+1} 5^l = \frac{1}{32} \left\{ \binom{6}{1} + \binom{6}{3} \times 5 + \binom{6}{5} \times 5^2 \right\} = \frac{1}{32}(6+100+150) = \frac{256}{32} = 8.$$

Property 2.2.

$$F_{k+2} = 1 + \sum_{i=1}^{k} F_i.$$

Proof. We have

$$\begin{aligned} F_0 &= 0, \\ F_1 &= 1, \\ F_2 &= F_0 + F_1, \\ F_3 &= F_1 + F_2, \\ &\vdots \\ F_{k+2} &= F_k + F_{k+1}. \end{aligned}$$

Adding these equalities we get

$$F_{k+2} = 1 + F_1 + F_2 + \ldots + F_k = 1 + \sum_{i=1}^{k} F_i.$$

□

Theorem 2.3.

$$F_{k+l} = F_l F_{k+1} + F_{l-1} F_k$$

with $k \in \mathbb{N}$ *and* $l \geq 2$. *(*$l \geq 2$*).*

Proof. We prove the result by induction on l.

From the definition of the Fibonacci sequence, we konw that $F_{k+2} = F_{k+1} + F_k = F_2 F_{k+1} + F_1 F_k$ since $F_1 = F_2 = 1$.

Let assume that: $F_{k+i} = F_i F_{k+1} + F_{i-1} F_k$ with $i = 2, 3, \ldots, l-1, l$.

Then, we have

$$F_{k+l+1} = F_{k+l} + F_{k+l-1}.$$

From our assumption, we have

$$F_{k+l} = F_l F_{k+1} + F_{l-1} F_k$$

and

$$F_{k+l-1} = F_{l-1} F_{k+1} + F_{l-2} F_k.$$

Thus we get

$$F_{k+l+1} = F_l F_{k+1} + F_{l-1} F_k + F_{l-1} F_{k+1} + F_{l-2} F_k,$$
$$F_{k+l+1} = (F_l + F_{l-1}) F_{k+1} + (F_{l-1} + F_{l-2}) F_k.$$

Using the recurrence relation of the Fibonacci sequence, we obtain

$$F_{k+l+1} = F_{l+1} F_{k+1} + F_l F_k.$$

□

Property 2.4. *Let* m *be a positive integer which is greater than* 2. *Then, we have*

$$F_{3m+2} = 4F_{3m-1} + F_{3m-4}.$$

Proof. For $m \geq 2$, we have

$$F_{3m+2} = F_{3m} + F_{3m+1} = F_{3m} + F_{3m} + F_{3m-1} = 2F_{3m} + F_{3m-1}$$
$$F_{3m+2} = 2(F_{3m-1} + F_{3m-2}) + F_{3m-1} = 3F_{3m-1} + 2F_{3m-2}$$
$$F_{3m+2} = 3F_{3m-1} + 2(F_{3m-3} + F_{3m-4}) = 3F_{3m-1} + 2F_{3m-3} + 2F_{3m-4}.$$

Again

$$F_{3m-1} = F_{3m-2} + F_{3m-3} = 2F_{3m-3} + F_{3m-4},$$

which gives

$$F_{3m-3} = \frac{1}{2}(F_{3m-1} - F_{3m-4}).$$

Thus

$$F_{3m+2} = 3F_{3m-1} + 2 \times \frac{1}{2}(F_{3m-1} - F_{3m-4}) + 2F_{3m-4}$$
$$F_{3m+2} = 3F_{3m-1} + F_{3m-1} - F_{3m-4} + 2F_{3m-4}$$
$$F_{3m+2} = 4F_{3m-1} + F_{3m-4}.$$

□

The above can be generalized to

Property 2.5. *Let* m *be a positive integer which is greater than* 2. *Then, we have*

$$F_{3m+2} = 4\sum_{i=2}^{m} F_{3i-1}.$$

Proof. From Property 2.4, we have for $m \geq 2$

$$\begin{aligned} F_2 &= 1, \\ F_5 &= 5, \\ F_8 &= 4F_5 + F_2, \\ F_{11} &= 4F_8 + F_5, \\ &\vdots \\ F_{3m+2} &= 4F_{3m-1} + F_{3m-4}. \end{aligned}$$

Adding the above gives the desired result. □

3. PROPERTIES OF THE FIBONACCI NUMBERS

Property 3.1. *$F_n \equiv 0 \pmod 2$ if and only if $n \equiv 0 \pmod 3$.*

Proof. We prove by induction that for $k \in \mathbb{N}$ we have

$$F_{3k} \equiv 0 \pmod 2$$

We have $F_0 = 0 \equiv 0 \pmod 2$.

We assume that $F_{3k} \equiv 0 \pmod 2$.

From Theorem 2.3 we have

$$F_{3(k+1)} = F_{3k+3} = F_3 F_{3k+1} + F_2 F_{3k}.$$

Since $F_3 = 2 \equiv 0 \pmod 2$ and we assumed that $F_{3k} \equiv 0 \pmod 2$, we have

$$F_{3(k+1)} \equiv 0 \pmod 2.$$

We again prove by induction for $k \in \mathbb{N}$

$$F_{3k+1} \equiv 1 \pmod 2.$$

We have $F_1 = 1 \equiv 1 \pmod 2$.

Let us assume that $F_{3k+1} \equiv 1 \pmod 2$.

From Theorem 2.3 we have

$$F_{3(k+1)+1} = F_{3k+4} = F_4 F_{3k+1} + F_3 F_{3k}.$$

Since $F_{3k} \equiv 0 \pmod 2$ and $F_4 = 3 \equiv 1 \pmod 2$, using the assumption $F_{3k+1} \equiv 1 \pmod 2$, it follows that

$$F_{3(k+1)+1} \equiv 1 \pmod 2.$$

From $F_{3k} \equiv 0 \pmod 2$ and $F_{3k+1} \equiv 1 \pmod 2$, using the recurence relation of the Fibonacci sequence, we also have

$$F_{3k+2} = F_{3k+1} + F_{3k} \equiv 1 \pmod 2.$$

Thus the property is established. □

Corollary 3.2. *If $p = 5k + 2$ is a prime which is strictly greater than 5 ($k \in \mathbb{N}$ and k odd), then $F_p = F_{5k+2}$ is an odd number.*

In order to prove this assertion, it suffices to remark that p is not divisible by 4.

Property 3.3.

$$F_{5k} \equiv 0 \pmod 5$$

with $k \in \mathbb{N}$.

Proof. We again prove this result by induction.

We have $F_0 = 0 \equiv 0 \pmod{5}$.

Notice also that $F_5 = 5 \equiv 0 \pmod{5}$.

Let assume that $F_{5k} \equiv 0 \pmod{5}$.

From Theorem 2.3 we have

$$F_{5(k+1)} = F_{5k+5} = F_5 F_{5k+1} + F_4 F_{5k}.$$

Since $F_5 \equiv 0 \pmod{5}$ and we assumed that $F_{5k} \equiv 0 \pmod{5}$, we obtain

$$F_{5(k+1)} \equiv 0 \pmod{5}.$$

□

Property 3.4.

$$F_n \geq n$$

with $n \in \mathbb{N}$ *and* $n \geq 5$.

Proof. We prove this by induction.

We have $F_5 = 5 \geq 5$.

Let assume that $F_i \geq i$ with $i = 5, 6, \ldots, n$.

From Property 2.2 using the assumption above, we have

$$F_{n+1} = 1 + \sum_{i=0}^{n-1} F_i \geq 1 + \sum_{i=0}^{n-1} i.$$

$$F_{n+1} \geq 1 + \frac{n(n-1)}{2}.$$

Again

$$\frac{n(n-1)}{2} - n = \frac{n(n-3)}{2} > 0.$$

So, from $F_{n+1} \geq 1 + \frac{n(n-1)}{2}$, we obtain

$$F_{n+1} \geq n + 1.$$

□

4. CHARACTERIZATION OF NUMBERS $5k + 2$ WHICH ARE PRIMES WITH $k \in \mathbb{N}$ AND k ODD

Let $p = 5k + 2$ be a prime number with k a non-zero positive integer which is odd. Notice that in this case, $5k \pm 1$ is an even number and so:

$$\left\lfloor \frac{5k \pm 1}{2} \right\rfloor = \frac{5k \pm 1}{2}.$$

Property 4.1.

$$F_{5k+2} \equiv 5k + 1 \pmod{5k+2}$$

with $k \in \mathbb{N}$ *and* k *odd such that* $5k + 2$ *is prime.*

Proof. From Theorems 1.8 and 2.1 we have

$$2^{5k+1} F_{5k+2} = \sum_{l=0}^{\frac{5k+1}{2}} \binom{5k+2}{2l+1} 5^l \equiv 5^{\frac{5k+1}{2}} \equiv 5k + 1 \pmod{5k+2}$$

where we used the fact that $\binom{5k+2}{2l+1}$ is divisible by $5k + 2$ for $l = 0, 1, \ldots, \frac{5k-1}{2}$.

From Fermat's little theorem, we have

$$2^{5k+1} \equiv 1 \pmod{5k+2}.$$

We get $F_{5k+2} \equiv 5k+1 \pmod{5k+2}$. □

Property 4.2.

$$F_{5k+1} \equiv 1 \pmod{5k+2}$$

with $k \in \mathbb{N}$ and k odd such that $5k+2$ is prime.

Proof. From Theorem 2.1 and Property 1.9 we have

$$2^{5k} F_{5k+1} = \sum_{l=0}^{\lfloor \frac{5k}{2} \rfloor} \binom{5k+1}{2l+1} 5^l \equiv (5k+1) \sum_{l=0}^{\lfloor \frac{5k}{2} \rfloor} 5^l \pmod{5k+2}.$$

We have

$$\sum_{l=0}^{\lfloor \frac{5k}{2} \rfloor} 5^l = \frac{5^{\lfloor \frac{5k}{2} \rfloor + 1} - 1}{4}.$$

We get from the above

$$2^{5k+2} F_{5k+1} \equiv (5k+1) \left\{ 5^{\lfloor \frac{5k}{2} \rfloor + 1} - 1 \right\} \pmod{5k+2}.$$

Since k is an odd positive integer, there exists a positive integer m such that $k = 2m+1$. It follows that

$$\left\lfloor \frac{5k}{2} \right\rfloor = 5m+2.$$

Notice that $5k+2 = 10m+7$ is prime, implies that $k \neq 5$ and $k \neq 11$ or equivalently $m \neq 2$ and $m \neq 5$. Other restrictions on k and m can be given.

From Theorem 1.8 we have

$$5^{5m+3} \equiv 10m+6 \pmod{10m+7}.$$

We can rewritte $\sum_{l=0}^{\lfloor \frac{5k}{2} \rfloor} 5^l = \frac{5^{\lfloor \frac{5k}{2} \rfloor + 1} - 1}{4}$ as

$$\sum_{l=0}^{\lfloor \frac{5k}{2} \rfloor} 5^l = \frac{5^{5m+3} - 1}{4}.$$

Moreover, we have

$$(5k+1) \left\{ 5^{\lfloor \frac{5k}{2} \rfloor + 1} - 1 \right\} = (10m+6) \left\{ 5^{5m+3} - 1 \right\}.$$

Or,

$$(10m+6) \left\{ 5^{5m+3} - 1 \right\} \equiv 5^{5m+3} \left\{ 5^{5m+3} - 1 \right\} \equiv 5^{10m+6} - 10m - 6 \pmod{10m+7}.$$

We have

$$(10m+6) \left\{ 5^{5m+3} - 1 \right\} \equiv 5^{10m+6} + 1 \pmod{10m+7}.$$

From Fermat's little theorem, we have $5^{10m+6} \equiv 1 \pmod{10m+7}$. Therefore

$$(10m+6) \left\{ 5^{5m+3} - 1 \right\} \equiv 2 \pmod{10m+7},$$

or equivalently

$$(5k+1) \left\{ 5^{\lfloor \frac{5k}{2} \rfloor + 1} - 1 \right\} \equiv 2 \pmod{5k+2}.$$

It follows that

$$2^{5k+2}F_{5k+1} \equiv 2 \pmod{5k+2}.$$

Since 2 and $5k+2$ are relatively prime

$$2^{5k+1}F_{5k+1} \equiv 1 \pmod{5k+2}.$$

From Fermat's little theorem, we have $2^{5k+1} \equiv 1 \pmod{5k+2}$. Therefore

$$F_{5k+1} \equiv 1 \pmod{5k+2}.$$

$\square$

Property 4.3.

$$F_{5k} \equiv 5k \pmod{5k+2}$$

with $k \in \mathbb{N}$ and k odd such that $5k+2$ is prime.

Proof. From Theorem 2.1 and Property 1.10 we have

$$2^{5k-1}F_{5k} = \sum_{l=0}^{\frac{5k-1}{2}} \binom{5k}{2l+1} 5^l \equiv \sum_{l=0}^{\frac{5k-1}{2}} (5k-2l)5^l \pmod{5k+2}.$$

Also

$$\sum_{l=0}^{\frac{5k-1}{2}} (5k-2l)5^l = \frac{5\left[3 \times 5^{\frac{5k-1}{2}} - (2k+1)\right]}{8}.$$

So

$$2^{5k+2}F_{5k} \equiv 5\left(3 \times 5^{\frac{5k-1}{2}} - (2k+1)\right) \pmod{5k+2}.$$

Moreover since $k = 2m+1$, we have

$$3 \times 5^{\frac{5k-1}{2}} - (2k+1) = 3 \times 5^{5m+2} - (4m+3).$$

Since $5^{5m+3} \equiv 10m+6 \pmod{10m+7}$, we have

$$3 \times 5^{5m+3} \equiv 30m+18 \pmod{10m+7},$$

$$3 \times 5^{5m+3} \equiv 40m+25 \pmod{10m+7}.$$

Consequently

$$3 \times 5^{5m+3} \equiv 8m+5 \pmod{10m+7}$$

which implies

$$3 \times 5^{5m+2} - (4m+3) \equiv 4m+2 \pmod{10m+7},$$

or equivalently for $k = 2m+1$

$$3 \times 5^{\frac{5k-1}{2}} - (2k+1) \equiv 2k \pmod{5k+2},$$

$$2^{5k+2}F_{5k} \equiv 2 \times 5k \pmod{5k+2}.$$

Since 2 and $5k+2$ are relatively prime

$$2^{5k+1}F_{5k} \equiv 5k \pmod{5k+2}.$$

From Fermat's little theorem, we have $2^{5k+1} \equiv 1 \pmod{5k+2}$. Therefore

$$F_{5k} \equiv 5k \pmod{5k+2}$$

$\square$

Property 4.4. *Let $5k+2$ be a prime with k odd and let m be a positive integer which is greater than 2. Then, we have*

$$F_{3m} \equiv 2\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i} F_{3i-1}\right) \pmod{5k+2}.$$

Proof. We prove the result by induction.

We know that $F_6 = 8$. Or, $2(3+F_2) = 2(3+1) = 2 \times 4 = 8$. So

$$F_6 = F_{3\times 2} = 2(3+F_2) \equiv 2(3+F_2) \pmod{5k+2}.$$

Let us assume that $F_{3m} \equiv 2\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i} F_{3i-1}\right) \pmod{5k+2}$ with $m \geq 2$.

For m a positive integer, we have by Theorem 2.3

$$F_{3(m+1)} = F_{3m+3} = F_3F_{3m+1} + F_2F_{3m} = 2F_{3m+1} + F_{3m}$$
$$F_{3(m+1)} = 2(F_{3m} + F_{3m-1}) + F_{3m} = 2F_{3m-1} + 3F_{3m}.$$

From the assumption above, we get

$$F_{3(m+1)} \equiv 2F_{3m-1} + 2\left(3^{m} + \sum_{i=1}^{m-1} 3^{m-i} F_{3i-1}\right) \pmod{5k+2},$$

$$F_{3(m+1)} \equiv 2\left(3^{m} + \sum_{i=1}^{m} 3^{m-i} F_{3i-1}\right) \pmod{5k+2}.$$

Thus the proof is complete by induction.

□

Theorem 4.5. *Let $5k+2$ be a prime with k an odd integer and let m be a positive integer which is greater than 2. Then*

$$F_{5mk} \equiv 5k\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i} F_{3i-1}\right) \pmod{5k+2}$$

and

$$F_{5mk+1} \equiv F_{3m-1} \pmod{5k+2}.$$

Proof. We prove the theorem by induction.

We have using Theorem 2.3

$$F_{10k} = F_{5k+5k} = F_{5k}F_{5k+1} + F_{5k-1}F_{5k} = F_{5k}(F_{5k+1} + F_{5k-1}).$$

Using Properties 4.1, 4.2 and 4.3 we can see that

$$F_{10k} \equiv 20k \pmod{5k+2}.$$

Also

$$5k\left(3 + \sum_{i=1}^{1} 3^{1-i} F_{3i-1}\right) = 5k(3+F_2) = 20k \equiv 20k \pmod{5k+2}.$$

So

$$F_{10k} = F_{5\times 2k} \equiv 5k\left(3 + \sum_{i=1}^{1} 3^{1-i} F_{3i-1}\right) \pmod{5k+2}.$$

Moreover, we have from Theorem 2.3, Property 4.2 and Property 4.3

$$F_{10k+1} = F_{5k+5k+1} = F_{5k+1}^2 + F_{5k}^2 \equiv 1 + 25k^2 \pmod{5k+2}.$$

We have

$$(5k+2)^2 = 25k^2 + 20k + 4 \equiv 25k^2 + 10k \equiv 0 \pmod{5k+2}.$$

So

$$25k^2 \equiv -10k \equiv 2(5k+2) - 10k \equiv 4 \pmod{5k+2}.$$

Therefore

$$F_{10k+1} \equiv F_5 \equiv 5 \pmod{5k+2}$$

or equivalently

$$F_{5\times 2k+1} \equiv F_{3\times 2-1} \equiv 5 \pmod{5k+2}.$$

Let us assume that

$$F_{5mk} \equiv 5k\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i} F_{3i-1}\right) \pmod{5k+2}$$

and

$$F_{5mk+1} \equiv F_{3m-1} \pmod{5k+2}$$

. Then, we have

$$F_{5(m+1)k} = F_{5mk+5k} = F_{5k}F_{5mk+1} + F_{5k-1}F_{5mk}.$$

Using Property 4.3 and $F_{5k-1} \equiv 3 \pmod{5k+2}$, from the assumptions above, we have

$$F_{5(m+1)k} \equiv 5kF_{3m-1} + 3 \times 5k\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i} F_{3i-1}\right) \pmod{5k+2}.$$

It gives

$$F_{5(m+1)k} \equiv 5k\left(3^{m} + \sum_{i=1}^{m} 3^{m-i} F_{3i-1}\right) \pmod{5k+2}.$$

Moreover, we have

$$F_{5(m+1)k+1} = F_{5mk+5k+1} = F_{5k+1}F_{5mk+1} + F_{5k}F_{5mk}.$$

Using Properties 4.2 and 4.3 and the assumptions above, and since $25k^2 \equiv 4 \pmod{5k+2}$, we have

$$F_{5(m+1)k+1} \equiv F_{3m-1} + 25k^2\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-i-1} F_{3i-1}\right) \pmod{5k+2},$$

$$F_{5(m+1)k+1} \equiv F_{3m-1} + 4\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-i-1} F_{3i-1}\right) \pmod{5k+2},$$

$$F_{5(m+1)k+1} \equiv F_{3m-1} + 2F_{3m} \pmod{5k+2}.$$

Or,

$$F_{3m-1} + 2F_{3m} = F_{3m-1} + F_{3m} + F_{3m} = F_{3m+1} + F_{3m} = F_{3m+2}.$$

Therefore

$$F_{5(m+1)k+1} \equiv F_{3m+2} \pmod{5k+2},$$

or equivalently

$$F_{5(m+1)k+1} \equiv F_{3(m+1)-1} \pmod{5k+2}.$$

This completes the proof. □

Corollary 4.6. *Let $5k+2$ be a prime with k an odd integer and m a positive integer which is greater than 2. Then*

$$F_{5mk+2} \equiv F_{3m-1} + 5k\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i}F_{3i-1}\right) \pmod{5k+2},$$

$$F_{5mk+3} \equiv 2F_{3m-1} + 5k\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i}F_{3i-1}\right) \pmod{5k+2},$$

and

$$F_{5mk+4} \equiv 3F_{3m-1} + 10k\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i}F_{3i-1}\right) \pmod{5k+2}.$$

Theorem 4.7. *Let $5k+2$ be a prime with k an odd positive integer, m a positive integer which is greater than 2 and $r \in \mathbb{N}$. Then*

$$F_{m(5k+r)} \equiv F_{mr}F_{3m-1} + 5kF_{mr-1}\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i}F_{3i-1}\right) \pmod{5k+2}.$$

Proof. For m, r two non-zero positive integers, we have by Theorem 2.3

$$F_{m(5k+r)} = F_{5mk+mr} = F_{mr}F_{5mk+1} + F_{mr-1}F_{5mk}.$$

From Theorem 4.5, we have for $m \geq 2$ and $r \in \mathbb{N}$

$$F_{m(5k+r)} \equiv F_{mr}F_{3m-1} + 5kF_{mr-1}\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i}F_{3i-1}\right) \pmod{5k+2}.$$

□

Remark 4.8. *In particular, if $r = 3$, we know that $F_{m(5k+3)} \equiv 0 \pmod{5k+2}$.*

This congruence can be deduced from Property 4.4 and Theorem 4.7. Indeed, using Theorem 4.7, we have

$$F_{m(5k+3)} \equiv F_{3m}F_{3m-1} + 5kF_{3m-1}\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i}F_{3i-1}\right) \pmod{5k+2},$$

$$F_{m(5k+3)} \equiv F_{3m}F_{3m-1} + 5kF_{3m-1}\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i}F_{3i-1}\right)$$
$$-(5k+2)F_{3m-1}\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i}F_{3i-1}\right) \pmod{5k+2},$$

$$F_{m(5k+3)} \equiv F_{3m}F_{3m-1} - 2F_{3m-1}\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i}F_{3i-1}\right) \pmod{5k+2}.$$

Using Property 4.4 we get

$$F_{m(5k+3)} \equiv F_{3m}F_{3m-1} - F_{3m-1}F_{3m} \equiv 0 \pmod{5k+2}.$$

Corollary 4.9. *Let $5k+2$ be a prime with k an odd positive integer and $m, r \in \mathbb{N}$. Then*

$$F_{m(5k+r)} \equiv F_{mr}F_{3m-1} - F_{mr-1}F_{3m} \pmod{5k+2}.$$

Lemma 4.10. *Let* $5k+2$ *be a prime with* k *an odd positive integer and* $r \in \mathbb{N}$*. Then*

$$F_{5k+r} \equiv F_r - 2F_{r-1} \pmod{5k+2}.$$

Proof. Let us prove the lemma by induction.

For $r = 1$, we have $F_{5k+r} = F_{5k+1} \equiv 1 \pmod{5k+2}$ and $F_r - 2F_{r-1} = F_1 - 2F_0 = F_1 \equiv 1 \pmod{5k+2}$.

Let us assume that $F_{5k+s} \equiv F_s - 2F_{s-1} \pmod{5k+2}$ for $s \in [[2, r]]$ with $r \geq 2$. Using the assumption, we have for $r \geq 2$

$$F_{5k+r+1} \equiv F_{5k+r} + F_{5k+r-1} \pmod{5k+2},$$

$$F_{5k+r+1} \equiv F_r - 2F_{r-1} + F_{r-1} - 2F_{r-2} \pmod{5k+2},$$

$$F_{5k+r+1} \equiv F_r + F_{r-1} - 2(F_{r-1} + F_{r-2}) \pmod{5k+2},$$

$$F_{5k+r+1} \equiv F_{r+1} - 2F_r \pmod{5k+2}.$$

Thus the lemma is proved. □

We now prove Corollary 4.9 as a consequence of Lemma 4.10.

Proof. For $m = 1$, we have $F_{mr}F_{3m-1} - F_{mr-1}F_{3m} = F_rF_2 - F_{r-1}F_3 = F_r - 2F_{r-1} \equiv F_{5k+r} \pmod{5k+2}$.

So, it remains to prove that for $m \geq 2$ and $r \in \mathbb{N}$, we have $F_{m(5k+r)} \equiv F_{mr}F_{3m-1} - F_{mr-1}F_{3m} \pmod{5k+2}$.

For $m \geq 2$ and $r \in \mathbb{N}$, from Theorem 4.7, we have

$$F_{m(5k+r)} \equiv F_{mr}F_{3m-1} + 5kF_{mr-1}\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i}F_{3i-1}\right) \pmod{5k+2},$$

$$F_{m(5k+r)} \equiv F_{mr}F_{3m-1} + 5kF_{mr-1}\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i}F_{3i-1}\right)$$

$$-(5k+2)F_{mr-1}\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i}F_{3i-1}\right) \pmod{5k+2},$$

$$F_{m(5k+r)} \equiv F_{mr}F_{3m-1} - 2F_{mr-1}\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i}F_{3i-1}\right) \pmod{5k+2}.$$

So, using Property 4.4, we deduce that for $m \geq 2$ and $r \in \mathbb{N}$

$$F_{m(5k+r)} \equiv F_{mr}F_{3m-1} - F_{mr-1}F_{3m} \pmod{5k+2}.$$

□

Corollary 4.11. *Let* $5k+2$ *be a prime with* k *an odd positive integer, let* m *be a positive integer and* $r \in \mathbb{N}$*. Then*

$$F_{m(5k+r)} \equiv F_{mr}F_{3m+1} - F_{mr+1}F_{3m} \pmod{5k+2}.$$

Lemma 4.12. *Let* $5k+2$ *be a prime with* k *an odd positive integer and let* m *be a positive integer. Then*

$$F_{5mk} + F_{3m} \equiv 0 \pmod{5k+2}.$$

Proof. For $m = 0$, we have $F_{5mk} + F_{3m} = 2F_0 = 0 \equiv 0 \pmod{5k+2}$.

For $m = 1$, we have $F_{5mk} + F_{3m} = F_{5k} + F_3 \equiv 5k + 2 \equiv 0 \pmod{5k+2}$.

So, it remains to prove that for $m \geq 2$, we have $F_{5mk} + F_{3m} \equiv 0 \pmod{5k+2}$.

From Theorem 4.5, we have

$$F_{5mk} \equiv 5k\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i} F_{3i-1}\right) \pmod{5k+2},$$

$$F_{5mk} \equiv 5k\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i} F_{3i-1}\right)$$

$$-(5k+2)\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i} F_{3i-1}\right) \pmod{5k+2},$$

$$F_{5mk} \equiv -2\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i} F_{3i-1}\right) \pmod{5k+2}.$$

From Property 4.4, we have $F_{5mk} \equiv -F_{3m} \pmod{5k+2}$. □

We now prove Corollary 4.11 as a consequence of Lemma 4.12

Proof. If $r = 0$, we can remark that $F_r F_{3m+1} - F_{r+1} F_{3m} = F_0 F_{3m+1} - F_1 F_{3m} = -F_{3m} \equiv F_{5mk} \pmod{5k+2}$.

Thus, it remains to prove that for $m \geq 2$ and $r \in \mathbb{N}$, $F_{m(5k+r)} \equiv F_{mr} F_{3m+1} - F_{mr+1} F_{3m} \pmod{5k+2}$.

From Corollary 4.9, we know that for $m, r \in \mathbb{N}$

$$F_{m(5k+r)} \equiv F_{mr} F_{3m-1} - F_{mr-1} F_{3m} \pmod{5k+2}.$$

Also

$$F_{mr-1} = F_{mr+1} - F_{mr}.$$

It follows that

$$F_{m(5k+r)} \equiv F_{mr} F_{3m-1} - F_{mr+1} F_{3m} + F_{mr} F_{3m} \pmod{5k+2},$$

$$F_{m(5k+r)} \equiv F_{mr}(F_{3m-1} + F_{3m}) - F_{mr+1} F_{3m} \pmod{5k+2},$$

$$F_{m(5k+r)} \equiv F_{mr} F_{3m+1} - F_{mr+1} F_{3m} \pmod{5k+2}.$$

□

Remark 4.13. *We can observe that* $F_{1\times(5k+r)} = F_{5k+r}$ *and* $F_{1\times r} F_{3\times 1+1} - F_{1\times r+1} F_{3\times 1} = F_r F_4 - F_{r+1} F_3 = 3F_r - 2F_{r+1}$. *By Properties 4.1, 4.2 and 4.3 we have*

$$\begin{aligned}
r = 1: \ & F_{5k+r} = F_{5k+1} \equiv 1 \pmod{5k+2},\ 3F_r - 2F_{r+1} = 3F_1 - 2F_2 = 1 \equiv 1 \pmod{5k+2}\\
r = 2: \ & F_{5k+r} = F_{5k+2} \equiv 5k+1 \pmod{5k+2},\ 3F_r - 2F_{r+1} = 3F_2 - 2F_3 = -1 \equiv 5k+1 \pmod{5k+2}\\
r = 3: \ & F_{5k+r} = F_{5k+3} \equiv 0 \pmod{5k+2},\ 3F_r - 2F_{r+1} = 3F_3 - 2F_4 = 0 \equiv 0 \pmod{5k+2}\\
r = 4: \ & F_{5k+r} = F_{5k+4} \equiv 5k+1 \pmod{5k+2},\ 3F_r - 2F_{r+1} = 3F_4 - 2F_5 = -1 \equiv 5k+1 \pmod{5k+2}
\end{aligned}$$

So, we have $F_{5k+r} \equiv 3F_r - 2F_{r+1} \pmod{5k+2}$ *or equivalently* $F_{1\times(5k+r)} \equiv F_{1\times r} F_{3\times 1+1} - F_{1\times r+1} F_{3\times 1} \pmod{5k+2}$ *with* $r \in [[1,4]]$.

Thus we have

Property 4.14. *Let $5k+2$ be a prime with k an odd positive integer, let m be a positive integer and $r \in \mathbb{N}$. Then*

$$F_{5k+r} \equiv 3F_r - 2F_{r+1} \pmod{5k+2}.$$

Proof. We have $F_{5k+0} = F_{5k} \equiv 5k \pmod{5k+2}$ and $3F_0 - 2F_1 = -2 \equiv 5k \pmod{5k+2}$. So, $F_{5k} \equiv 3F_0 - 2F_1 \pmod{5k+2}$.

Moreover, we know that $F_{5k+1} \equiv 3F_1 - 2F_2 \equiv 1 \pmod{5k+2}$.

Let us assume that $F_{5k+s} \equiv 3F_s - 2F_{s+1} \pmod{5k+2}$ for $s \in [[1, r]]$. We have for $r \in \mathbb{N}$,

$$F_{5k+r+1} = F_{5k+r} + F_{5k+r-1} \equiv 3F_r - 2F_{r+1} + 3F_{r-1} - 2F_r \pmod{5k+2}$$

$$F_{5k+r+1} \equiv 3(F_r + F_{r-1}) - 2(F_{r+1} + F_r) \pmod{5k+2}$$

$$F_{5k+r+1} \equiv 3F_{r+1} - 2F_{r+2} \pmod{5k+2}.$$

Thus the proof is complete by induction. ☐

Property 4.15. *Let $5k+2$ be a prime with k odd and let m be a positive integer which is greater than 2. Then, we have*

$$F_{3m+1} = 3^m + 2\sum_{i=1}^{m-1} 3^{m-1-i} F_{3i} \pmod{5k+2}.$$

Proof. We prove the result by induction.

We have for $m = 2$, $F_{3m+1} = F_{3\times 2+1} = F_7 = 13$ and $3^m + 2\sum_{i=1}^{m-1} 3^{m-1-i} F_{3i} = 3^2 + 2F_3 = 9 + 2 \times 2 = 9 + 4 = 13$. So, $F_7 \equiv 3^2 + 2F_3 \equiv 13 \pmod{5k+2}$.

Let us assume that for $m \geq 2$, $F_{3m+1} = 3^m + 2\sum_{i=1}^{m-1} 3^{m-1-i} F_{3i} \pmod{5k+2}$. Using this assumption, we have for $m \geq 2$

$$F_{3(m+1)+1} = F_{3m+4} = F_4 F_{3m+1} + F_3 F_{3m} = 3F_{3m+1} + 2F_{3m}$$

$$F_{3(m+1)+1} \equiv 3^{m+1} + 2\sum_{i=1}^{m-1} 3^{m-i} F_{3i} + 2F_{3m} \pmod{5k+2},$$

$$F_{3(m+1)+1} \equiv 3^{m+1} + 2\sum_{i=1}^{m} 3^{m-i} F_{3i} \pmod{5k+2}.$$

Thus the induction hypothesis holds. ☐

Corollary 4.16. *Let $5k+2$ be a prime with k odd and let m be a positive integer which is greater than 2. Then, we have*

$$F_{3m+2} = 3^m + 2\left(3^{m-1} + \sum_{i=1}^{m-1} 3^{m-1-i} F_{3i+1}\right) \pmod{5k+2}.$$

Proof. It stems from the recurrence relation of the Fibonacci sequence which implies that $F_{3m+2} = F_{3m} + F_{3m+1}$ and $F_{3k+1} = F_{3k} + F_{3k-1}$ and Properties 4.4 and 4.15. ☐

5. PERIODS OF THE FIBONACCI SEQUENCE MODULO A POSITIVE INTEGER

Notice that $F_1 = F_2 \equiv 1 \pmod{m}$ with m an integer which is greater than 2.

Definition 5.1. *The Fibonacci sequence* (F_n) *is periodic modulo a positive integer* m *which is greater than* 2 $(m \geq 2)$*, if there exists at least a non-zero integer* ℓ_m *such that:*

$$F_{1+\ell_m} \equiv F_{2+\ell_m} \equiv 1 \pmod{m}$$

The number ℓ_m *is called a period of the Fibonacci sequence* (F_n) *modulo* m.

Remark 5.2. *For* $m \geq 2$ *we have* $l_m \geq 2$. *Indeed,* ℓ_m *cannot be equal to* 1 *since* $F_3 = 2$.

From Theorem 2.3 we have

$$F_{2+\ell_m} = F_{\ell_m} F_3 + F_{\ell_m - 1} F_2 \equiv 2F_{\ell_m} + F_{\ell_m - 1} \pmod{m}.$$

Since $F_{\ell_m} + F_{\ell_m-1} = F_{1+\ell_m}$, we get

$$F_{2+\ell_m} \equiv 2F_{\ell_m} + F_{\ell_m-1} \equiv F_{\ell_m} + F_{1+\ell_m} \equiv F_{\ell_m} + F_{2+\ell_m} \pmod{m}.$$

Therefore we have the following

Property 5.3.

$$F_{\ell_m} \equiv 0 \pmod{m}$$

Moreover, from Theorem 2.3 we have

$$F_{1+\ell_m} = F_{\ell_m} F_2 + F_{\ell_m-1} F_1 \equiv F_{\ell_m} + F_{\ell_m-1} \equiv F_{\ell_m-1} \pmod{m}.$$

Since $F_{1+\ell_m} \equiv 1 \pmod{m}$, we obtain the following

Property 5.4.

$$F_{\ell_m-1} \equiv 1 \pmod{m}.$$

Besides, using the recurrence relation of the Fibonacci sequence, from Property 5.3 we get

$$F_{\ell_m-2} + F_{\ell_m-1} = F_{\ell_m} \equiv 0 \pmod{m}.$$

Using Property 5.4 we obtain

Property 5.5.

$$F_{\ell_m-2} \equiv m - 1 \pmod{m}.$$

Remark 5.6. *From Theorem 2.3 we have for* $m \geq 2$

$$F_{2m} = F_{m+m} = F_m F_{m+1} + F_{m-1} F_m = F_m (F_{m+1} + F_{m-1})$$

and

$$F_{2m+1} = F_{(m+1)+m} = F_m F_{m+2} + F_{m-1} F_{m+1} = F_m (F_m + F_{m+1}) + F_{m-1}(F_{m-1} + F_m),$$

$$F_{2m+1} = F_m (2F_m + F_{m-1}) + F_{m-1}(F_{m-1} + F_m) = 2F_m^2 + 2F_m F_{m-1} + F_{m-1}^2 = F_m^2 + F_{m+1}^2.$$

From this we get

$$F_{2m+2} = F_{2m+1} + F_{2m} = F_m^2 + F_{m+1}^2 + F_m(F_{m+1} + F_{m-1}),$$

$$F_{2m+3} = F_3 F_{2m+1} + F_2 F_{2m} = 2(F_m^2 + F_{m+1}^2) + F_m(F_{m+1} + F_{m-1}),$$

$$F_{2m+4} = F_{2m+3} + F_{2m+2} = 3(F_m^2 + F_{m+1}^2) + 2F_m(F_{m+1} + F_{m-1}).$$

Theorem 5.7. *A period of the Fibonacci sequence modulo* $5k + 2$ *with* $5k + 2$ *a prime and* k *odd is given by*

$$\ell_{5k+2} = 2(5k + 3).$$

Proof. Using the recurrence relation of the Fibonacci sequence, and from Properties 4.1, 4.2 and 4.3 we have

$$F_{5k+3} = F_{5k+2} + F_{5k+1} \equiv 5k+2 \equiv 0 \pmod{5k+2}.$$

Taking $m = 5k+2$ prime (k odd) in the formulas of F_{2m+3} and F_{2m+4}, we have

$$F_{10k+7} = 2(F_{5k+2}^2 + F_{5k+3}^2) + F_{5k+2}(F_{5k+3} + F_{5k+1}) \equiv 2(5k+1)^2 + 5k + 1 \pmod{5k+2},$$

$$F_{10k+7} \equiv 50k^2 + 20k + 2 + 5k + 1 \equiv 10k(5k+2) + (5k+2) + 1 \pmod{5k+2},$$

$$F_{10k+7} \equiv 1 + (10k+1)(5k+2) \equiv 1 \pmod{5k+2},$$

and

$$F_{10k+8} = 3(F_{5k+2}^2 + F_{5k+3}^2) + 2F_{5k+2}(F_{5k+3} + F_{5k+1}) \equiv 3(5k+1)^2 + 2(5k+1) \pmod{5k+2},$$

$$F_{10k+8} \equiv 75k^2 + 30k + 3 + 10k + 2 \equiv 15k(5k+2) + 2(5k+2) + 1 \pmod{5k+2},$$

$$F_{10k+8} \equiv 1 + (15k+2)(5k+2) \equiv 1 \pmod{5k+2}.$$

Thus

$$F_{10k+7} \equiv F_{10k+8} \equiv 1 \pmod{5k+2},$$

or equivalently

$$F_{1+2(5k+3)} \equiv F_{2+2(5k+3)} \equiv 1 \pmod{5k+2}.$$

We deduce that a period of the Fibonacci sequence modulo $5k+2$ with $5k+2$ a prime is $\ell_{5k+2} = 2(5k+3)$. □

Remark 5.8. *We can observe that*

$$F_{5k-1} = F_{5k+1} - F_{5k} \equiv 1 - 5k \equiv 3 \equiv F_4 \pmod{5k+2},$$

$$F_{5k-2} = F_{5k} - F_{5k-1} \equiv 5k - 3 \equiv 5k - F_4 \pmod{5k+2},$$

$$F_{5k-3} = F_{5k-1} - F_{5k-2} \equiv 6 - 5k \equiv 8 \equiv F_6 \pmod{5k+2},$$

$$F_{5k-4} = F_{5k-2} - F_{5k-3} \equiv 5k - 11 \equiv 5k - (F_4 + F_6) \pmod{5k+2}.$$

Using induction we can show the following two properties.

Property 5.9. *Let $5k+2$ be a prime with k odd. Then, we have*

$$F_{5k-(2l+1)} \equiv F_{2(l+2)} \pmod{5k+2}$$

with l a positive integer such that $l \leq \lfloor \frac{5k-1}{2} \rfloor$.

Property 5.10. *Let $5k+2$ be a prime with k odd. Then, we have*

$$F_{5k-2l} \equiv 5k - \sum_{i=0}^{l-1} F_{2(i+2)} \pmod{5k+2}$$

with $l \geq 1$ such that $l \leq \lfloor \frac{5k}{2} \rfloor$.

Remark 5.11. *We can notice that*

$$F_{5k+4} = F_{5k+3} + F_{5k+2} \equiv F_{5k+2} \equiv 5k+1 \pmod{5k+2},$$

$$F_{5k+5} = F_{5k+4} + F_{5k+3} \equiv F_{5k+4} \equiv 5k+1 \pmod{5k+2},$$

$$F_{5k+6} = F_{5k+5} + F_{5k+4} \equiv 10k+2 \equiv 5k \pmod{5k+2},$$

and for $l \geq 1$ *we have*

$$F_{5k+3l+2} = F_{3l+2}F_{5k+1} + F_{3l+1}F_{5k} \equiv F_{3l+2} + 5kF_{3l+1} \pmod{5k+2},$$

$$F_{5k+3l+2} \equiv F_{3l}+(5k+1)F_{3l+1} \equiv F_{3l}-F_{3l+1}+(5k+2)F_{3l+1} \equiv F_{3l}-F_{3l+1} \equiv -F_{3l-1} \pmod{5k+2}.$$

Furthermore, we have for $l \geq 1$

$$F_{5k+3l+1} = F_{3l+1}F_{5k+1} + F_{3l}F_{5k} \equiv F_{3l+1} + 5kF_{3l} \pmod{5k+2},$$

$$F_{5k+3l+1} \equiv F_{3l-1}+(5k+1)F_{3l} \equiv F_{3l-1}-F_{3l}+(5k+2)F_{3l} \equiv F_{3l-1}-F_{3l} \equiv -F_{3l-2} \pmod{5k+2}.$$

Besides, we have for $l \geq 1$

$$F_{5k+3l} = F_{3l}F_{5k+1} + F_{3l-1}F_{5k} \equiv F_{3l} + 5kF_{3l-1} \pmod{5k+2},$$

$$F_{5k+3l} \equiv F_{3l-2}+(5k+1)F_{3l-1} \equiv F_{3l-2}-F_{3l-1}+(5k+2)F_{3l-1} \equiv F_{3l-2}-F_{3l-1} \equiv -F_{3l-3} \pmod{5k+2}.$$

We can state the following property, the proof of which follows from the above remark and by using induction.

Property 5.12. $F_{5k+n} \equiv -F_{n-3} \pmod{5k+2}$.

Theorem 5.13. *Let* $5k+2$ *be a prime with* k *an odd positive number and let* n *a positive integer. Then, we have*

$$F_{n(5k+3)} \equiv 0 \pmod{5k+2}.$$

Proof. The proof of the theorem will be done by induction. We have $F_0 \equiv 0 \pmod{5k+2}$. Moreover, we know that $F_{5k+3} \equiv 0 \pmod{5k+2}$. Let us assume that $F_{n(5k+3)} \equiv 0 \pmod{5k+2}$. We have

$$F_{(n+1)(5k+3)} = F_{n(5k+3)+5k+3} = F_{5k+3}F_{n(5k+3)+1} + F_{5k+2}F_{n(5k+3)}.$$

Since $F_{5k+3} \equiv 0 \pmod{5k+2}$, using the assumption $F_{n(5k+3)} \equiv 0 \pmod{5k+2}$, we deduce that

$$F_{(n+1)(5k+3)} \equiv 0 \pmod{5k+2}.$$

□

The following follows very easily from the above theorem.

Corollary 5.14. *If* $5k+3|m$, *then* $F_m \equiv 0 \pmod{5k+2}$.

Remark 5.15. *From the definition of a period of the Fibonacci sequence modulo an integer which is greater than* 2, *if* $F_{m(5k+r)+1} \equiv F_{m(5k+r)+2} \equiv 1 \pmod{5k+2}$, *then the number* $m(5k+r)$ *is a period of the Fibonacci sequence modulo* $5k+2$ *and we have* $F_{m(5k+r)} \equiv 0 \pmod{5k+2}$.

Theorem 5.16. *Let $5k+2$ be a prime with k an odd positive integer and r a positive integer which is strictly greater than 3. If there exists an integer m which is strictly greater than 2 such that*

$$F_{m(5k+r)} \equiv 0 \pmod{5k+2}$$

and

$$F_{3m+1} \equiv F_{3m} \pmod{5k+2}$$

with F_{3m} and $5k+2$ relatively prime, then a period of the Fibonacci sequence can be written independently of r either as a multiple of $3m-1$, or as a multiple of $5k+3$ or as a multiple of $5mk+1$.

Proof. Since $F_{3m+1} \equiv F_{3m} \pmod{5k+2}$, then $F_{3m-1} = F_{3m+1} - F_{3m} \equiv 0 \pmod{5k+2}$. So, $3m-1$ with $m > 2$ may represent a period of the Fibonacci sequence modulo $5k+2$ with $5k+2$ prime (k odd).

Moreover, if there exists an integer $m > 2$ such that $F_{m(5k+r)} \equiv 0 \pmod{5k+2}$, then from Corollary 4.11, we have for $m > 2$ and $r > 3$

$$F_{mr}F_{3m+1} \equiv F_{mr+1}F_{3m} \pmod{5k+2}.$$

Since $F_{3m+1} \equiv F_{3m} \pmod{5k+2}$ with F_{3m} and $5k+2$ relatively prime, it gives for $m > 2$ and $r > 3$

$$F_{mr} \equiv F_{mr+1} \pmod{5k+2}.$$

The congruences $F_{3m+1} \equiv F_{3m} \pmod{5k+2}$ and $F_{mr} \equiv F_{mr+1} \pmod{5k+2}$ with $F_{m(5k+r)} \equiv 0 \pmod{5k+2}$ imply that

$$F_{m(5k+r)} \equiv F_{3m-1} \equiv F_{mr-1} \equiv 0 \pmod{5k+2}.$$

So, there exists a non-zero integer ℓ_1 which may represent a period of the Fibonacci sequence modulo $5k+2$ such that

$$mr-1 = 3m-1+\ell_1.$$

We verify that $F_{\ell_1} \equiv 0 \pmod{5k+2}$. Indeed, we have

$$F_{3m-1+\ell_1} = F_{\ell_1+3m-1} = F_{3m-1}F_{\ell_1+1} + F_{3m-2}F_{\ell_1}.$$

Also

$$F_{3m-2} = F_{3m} - F_{3m-1} \equiv F_{3m} \pmod{5k+2}.$$

Since $F_{3m-1} \equiv 0 \pmod{5k+2}$ and $F_{3m-1+\ell_1} = F_{mr-1} \equiv 0 \pmod{5k+2}$, it follows that

$$F_{3m-1+\ell_1} \equiv F_{3m}F_{\ell_1} \equiv 0 \pmod{5k+2}.$$

Since F_{3m} and $5k+2$ are relatively prime

$$F_{\ell_1} \equiv 0 \pmod{5k+2}.$$

From $mr-1 = 3m-1+\ell_1$, we obtain

$$\ell_1 = m(r-3).$$

Moreover, since $F_{\ell_1} \equiv 0 \pmod{5k+2}$, there exists a non-zero integer ℓ_2 which may represent a period of the Fibonacci sequence modulo $5k+2$ such that

$$m(5k+r) = \ell_1 + \ell_2 = m(r-3) + \ell_2$$

which gives

$$\ell_2 = m(5k+3).$$

Since $F_{m(5k+3)} \equiv 0 \pmod{5k+2}$, we verify that $F_{\ell_2} \equiv 0 \pmod{5k+2}$.

The case $m = 1$ is not possible. Otherwise, from the definition of a period of the Fibonacci sequence modulo an integer which is greater than 2, F_{5k+4} and F_{5k+5} would be congruent to 1 modulo $5k + 2$. Or, $F_{5k+4} \equiv F_{5k+5} \equiv 5k + 1 \pmod{5k + 2}$. So, we reach a contradiction.

We have shown that $2(5k + 3)$ is a period of the Fibonacci sequence modulo $5k + 2$. But, we assume that $m > 2$ and we cannot have $F_{3m} \equiv F_{3m+1} \pmod{5k + 2}$ when $m = 2$ (recall that $5k + 2$ with k odd, is prime).

Regarding the equation $F_{m(5k+r)} \equiv F_{mr-1} \equiv 0 \pmod{5k + 2}$, since $F_{\ell_1} \equiv 0 \pmod{5k + 2}$, we can see also that there exists a non-zero integer ℓ_3 which may represent a period of the Fibonacci sequence such that

$$m(5k + r) = mr - 1 + \ell_3$$

which gives

$$\ell_3 = 5mk + 1.$$

Since $F_{3m-1} \equiv 0 \pmod{5k + 2}$ when $F_{3m} \equiv F_{3m+1} \pmod{5k + 2}$, we verify that $F_{\ell_3} = F_{5mk+1} \equiv F_{3m-1} \equiv 0 \pmod{5k + 2}$. The cases $m = 1$ and $m = 2$ are not possible since $F_{5k+1} \equiv 1 \pmod{5k + 2}$ and $F_{10k+1} \equiv 5 \pmod{5k + 2}$.

Regarding the equation $F_{m(5k+r)} \equiv F_{3m-1} \equiv 0 \pmod{5k + 2}$, since $F_{\ell_1} \equiv 0 \pmod{5k + 2}$, there exists a non-zero integer ℓ_4 such that

$$m(5k + r) = 3m - 1 + \ell_4.$$

We verify that $F_{\ell_4} \equiv 0 \pmod{5k + 2}$. Indeed, we have

$$F_{3m-1+\ell_4} = F_{\ell_4+3m-1} = F_{3m-1}F_{\ell_4+1} + F_{3m-2}F_{\ell_4}.$$

And

$$F_{3m-1+\ell_4} \equiv F_{3m}F_{\ell_4} \equiv 0 \pmod{5k + 2}.$$

Since we assume that F_{3m} and $5k + 2$ are relatively prime, we get

$$F_{\ell_4} \equiv 0 \pmod{5k + 2}.$$

From $m(5k + r) = 3m - 1 + \ell_4$, we obtain

$$\ell_4 = m(5k + r - 3) + 1.$$

When $F_{m(5k+r)} \equiv 0 \pmod{5k+2}$ and $F_{3m+1} \equiv F_{3m} \pmod{5k+2}$ such that F_{3m} and $5k+2$ relatively prime with $5k + 2$ prime (k odd), $r > 3$ and $m > 2$, the numbers $3m - 1$, ℓ_1, ℓ_2, ℓ_3 and ℓ_4 represent all the possibilities to write a period of the Fibonacci sequence modulo $5k+2$ with $5k + 2$ prime (k odd).

Therefore, assuming that $F_{m(5k+r)} \equiv 0 \pmod{5k + 2}$ and $F_{3m+1} \equiv F_{3m} \pmod{5k + 2}$ with $m > 2$ and $r > 3$, a period of the Fibonacci sequence modulo $5k + 2$ with $5k + 2$ prime (k odd), can be written independently of r either as a multiple of $3m - 1$, either as a multiple of $5k + 3$ or as a multiple of $5mk + 1$. □

Corollary 5.17. *A period of the Fibonacci sequence modulo $5k + 2$ with $5k + 2$ prime and k odd can be written as a multiple of $5k + 3$.*

Proof. We can notice that (see the proof of Theorem 5.16 for the notations)

$$\ell_3 + 3m - 1 = m(5k + 3) = \ell_4.$$

It is clear that if $3m - 1$ and ℓ_3 are periods of the Fibonacci sequence modulo $5k + 2$ with $5k+2$ prime (k odd), then $\ell_4 = \ell_3 + 3m - 1$ is also a period of the Fibonacci sequence modulo $5k + 2$.

Moreover, it is alway possible to find non-zero positive integers m_i with $i = 0, 1, 2, 3, 4$ such that:

$$\ell_i = m_i(5k+3) = m_i(5k+2) + m_i \equiv m_i \pmod{5k+2}$$

where we set $\ell_0 = 3m - 1$.

We now prove the existence of such m_i, with $i = 0, 1, 2, 3, 4$.

Indeed, when $i = 0$, we have $m_0 \equiv 3m - 1 \pmod{5k+2}$. We set $m_0 + 1 = 3t_0$ with $3t_0 - 1 > 0$. Then, since 3 and $5k+2$ are relatively prime, $t_0 \equiv m \pmod{5k+2}$. So, there exists an integer s_0 such that $m = t_0 + s_0(5k+2)$. Since $m > 2$ and $t_0 > 0$, we have necessarily $t_0 > -s_0(5k+2)$, $m - s_0(5k+2) \geq 1$ and so $3(m - s_0(5k+2)) - 1 \geq 2$. It implies that $3m - 1 = 3t_0 - 1 + 3s_0(5k+2) = m_0 + 3s_0(5k+2)$. It gives

$$m_0 = 3(m - s_0(5k+2)) - 1 > 0.$$

When $i = 1$, we have $m_1 \equiv m(r-3) \pmod{5k+2}$. If $r - 3 \equiv 0 \pmod{5k+2}$, then $m_1 \equiv 0 \pmod{5k+2}$. In this case, we can find two non-zero positive integers u_1 and v_1 such that $m_1 = u_1(5k+2)$ and $r - 3 = v_1(5k+2)$. Since $m(r-3) = m_1(5k+3)$, it leads to $mv_1 = u_1(5k+3)$. Since $5k+3 | mv_1$ and so $5k+3 | m(r-3)$, then we have $u_1 = \frac{mv_1}{5k+3} \in \mathbb{N}$ and for $r > 3$

$$m_1 = \frac{mv_1(5k+2)}{5k+3} = \frac{m(r-3)}{5k+3} \in \mathbb{N}.$$

If $r - 3$ and $5k+2$ are relatively prime, we set $m_1 = t_1(r-3)$ with $t_1 > 0$. Then, since $r - 3$ and $5k+2$ are relatively prime, $t_1 \equiv m \pmod{5k+2}$. So, there exists an integer s_1 such that $m = t_1 + s_1(5k+2)$. Since $m > 2$ and $t_1 > 0$, we have necessarily $t_1 > -s_1(5k+2)$ and $m - s_1(5k+2) > 0$. It implies that $m(r-3) = t_1(r-3) + s_1(r-3)(5k+2) = m_1 + s_1(r-3)(5k+2)$. It gives for $r > 3$

$$m_1 = (r-3)(m - s_1(5k+2)) > 0.$$

When $i = 2$, we have $\ell_2 = m(5k+3) = m_2(5k+3)$. So

$$m_2 = m > 0.$$

When $i = 3$, we have $m_3 \equiv 5mk + 1 \pmod{5k+2}$. We set $m_3 - 1 = 5kt_3$ with $t_3 > 0$. Then, since $5k$ and $5k+2$ are relatively primes, $t_3 \equiv m \pmod{5k+2}$. So, there exists an integer s_3 such that $m = t_3 + s_3(5k+2)$. Since $m > 2$ and $t_3 > 0$, we have necessarily $t_3 > -s_3(5k+2)$ and $m - s_3(5k+2) > 0$. It implies that $5mk = 5kt_3 + 5ks_3(5k+2) = m_3 - 1 + 5ks_3(5k+2)$. It gives

$$m_3 = 5k(m - s_3(5k+2)) + 1 > 0.$$

When $i = 4$, we have $m_4 \equiv m(5k+r-3) + 1 \pmod{5k+2}$. If $5k+r-3 \equiv 0 \pmod{5k+2}$, then $m_4 \equiv 0 \pmod{5k+2}$. In this case, we can find two non-zero positive integers u_4 and v_4 such that $m_4 = u_4(5k+2)$ and $r - 3 = v_4(5k+2)$. Since $m(5k+r-3) = m_4(5k+3)$, it leads to $mv_4 = u_4(5k+3)$. Since $5k+3 | mv_4$ and so $5k+3 | m(5k+r-3)$, then we have $u_4 = \frac{mv_4}{5k+3} \in \mathbb{N}$ and for $r > 3$

$$m_4 = \frac{mv_4(5k+2)}{5k+3} = \frac{m(5k+r-3)}{5k+3} \in \mathbb{N}.$$

If $5k+r-3$ and $5k+2$ are relatively prime, we set $m_4 = t_4(5k+r-3)$ with $t_4 > 0$. Then, since $5k+r-3$ and $5k+2$ are relatively prime, $t_4 \equiv m \pmod{5k+2}$. So, there exists an integer s_4 such that $m = t_4 + s_4(5k+2)$. Since $m > 2$ and $t_4 > 0$, we have necessarily $t_4 > -s_4(5k+2)$

and $m-s_4(5k+2) > 0$. It implies that $m(5k+r-3) = t_4(5k+r-3)+s_4(5k+r-3)(5k+2) = m_4 + s_4(5k+r-3)(5k+2)$. It gives for $r > 3$

$$m_4 = (5k+r-3)(m-s_4(5k+2)) > 0.$$

Thus, the numbers m_i with $i = 0, 1, 2, 3, 4$ exist. Therefore, it is always possible to find an integer m_i with $i = 0, 1, 2, 3, 4$ such that $\ell_i = m_i(5k+3)$.

Consequently, any period of the Fibonacci sequence modulo $5k+2$ with $5k+2$ prime (k odd), can be written as a multiple of $5k+3$. □

Corollary 5.18. *The minimal period of the Fibonacci sequence modulo* $5k+2$ *with* $5k+2$ *prime and* k *odd is* $2(5k+3)$.

We know that $2(5k+3)$ is a period of the Fibonacci sequence modulo $5k+2$ with $5k+2$ prime with k odd. Moreover, $5k+3$ cannot be a period of the Fibonacci sequence modulo $5k+2$ with $5k+2$ prime and k odd since $F_{5k+4} \equiv F_{5k+5} \equiv 5k+1 \pmod{5k+2}$. So, Corollary 5.18 follows from Corollary 5.17.

6. Acknowledgements

The authors would like to thank Zakir Ahmed, Sudarson Nath and Ankush Goswami for helpful comments.

References

[1] D. M. Burton, *Elementary Number Theory*, Tata McGraw Hill, 2007.
[2] R. A. Mollin, *Fundamental Number Theory with Applications*, Chapman and Hall/CRC, 2008.

MSC2010: 11A07, 11B37, 11B39, 11B50.

Lycée professionnel hotelier La Closerie, 10 rue Pierre Loti - BP 4, 22410 Saint-Quay-Portrieux, France
E-mail address: `laugier.alexandre@orange.fr`

Department of Mathematical Sciences, Tezpur University, Napaam, Sonitpur, Assam, Pin 784028, India
E-mail address: `manjil@gonitsora.com`